

Hydrogen Abstraction Reactions in the Kiloelectronvolt Particle Bombardment of Organic Films

Ramona S. Taylor and Barbara J. Garrison*

Department of Chemistry
152 Davey Laboratory
The Pennsylvania State University
University Park, Pennsylvania 16802

Received September 27, 1993

The bombardment of molecular solids with high-energy (500–25 000 eV) particles is an effective method of placing these molecules, as both ions and neutrals, into the gas phase for subsequent analysis via mass spectrometry (MS) techniques such as secondary ion MS (SIMS),¹ fast atom bombardment MS (FAB),^{1e,h} and laser postionization MS.^{1a,b} In addition to that of the intact parent molecule, peaks corresponding to fragments of the molecule, the molecule plus or minus a proton, and even various aggregates of the molecule are often found in the mass spectrum.¹⁻⁴ A number of mechanisms have been proposed for the formation of these detected species. Clusters smaller than the parent molecule are believed to result either from the direct ejection of a portion of the intact molecule^{2,3,5,6} or from the unimolecular dissociation of larger transitory clusters during their flight to the detector.⁵ Clusters larger than the parent molecule are proposed to result from either the near-surface reaction of two previously ejected entities,⁴⁻⁷ the direct ejection of a large precursor ion created prior to the bombardment event,^{1d,1f} or an oxidation or reduction reaction among escaping molecules within the selva area, the solid-gas interfacial region most distorted by the bombarding particle.^{1d-f,8} In this communication, we propose a new mechanism for molecule formation during the kilo-electronvolt (keV) particle bombardment of an organic film. Molecular dynamics computer simulations using a many-body interaction potential to describe the hydrocarbon interactions show a novel H abstraction mechanism whereby a free H atom formed during the collision event within ~ 100 fs abstracts another H atom from a stationary neighboring surface molecule to form gaseous H₂. Recombination type mechanisms involving lone H atoms have previously been proposed as unlikely because hydrogen's light mass usually results in a high ejection velocity.⁹ Consequently, within a very short time, the atom is beyond the interaction range of the other heavier moving particles. The mechanism proposed here occurs on a short time scale and is a direct consequence of collisional motion. This is in contrast to previously proposed reactions for H transfer in which the reactions are extensions of thermal energy processes. The prediction here

that free H atoms can react with stationary molecules before leaving the interaction area suggests new directions for explaining chemical reactions in organic films exposed to keV particle bombardment.

Molecular dynamics (MD) simulations have been successful in yielding insight into ejection processes ranging from emission and angular distributions to mechanisms of molecular emission that occur in the keV particle bombardment of solids.^{5-7,9-14} Although previous MD simulations of the keV particle bombardment of organic films explained some of the ejection mechanisms in these systems, the potential energy functions used to model the adsorbates warranted only the exploration of molecular ejection and fragmentation, and not that of any chemical reactions which might occur among atoms originating from different adsorbate molecules.^{5,6} Recently, a many-body interaction potential for hydrocarbons was developed that describes reactions among all atoms including those from different adsorbate molecules.¹⁵ Thus, the possibility now exists to explore the importance of intermolecular reactions.

As a prototypical system for the keV bombardment of an organic film, the bombardment of a p(2 \times 2) overlayer of ethylidyne (C₂H₃) adsorbed on the {111} face of Pt with a 500-eV Ar beam was chosen for investigation. C₂H₃, a stable intermediate in the Pt-catalyzed dehydrogenation of ethylene and acetylene,^{2,16} adsorbs in a 3-fold hollow site through three Pt-C bonds with its C-C bond axis perpendicular to the Pt surface.¹⁶ The molecular dynamics scheme being used has been described in great detail elsewhere.^{10,11} Here, we include many-body potential energy functions which allow for chemical reactions among all of the constituents of the target. Thus, as the collision cascade evolves, reactions occurring among the substrate atoms, the substrate and adsorbate atoms, and the individual adsorbate atoms can be followed in time. The Pt-Pt interactions are described with a potential function based upon the embedded atom method (EAM).^{11,17} The C-C, C-H, and H-H interactions are described with a Tersoff-like potential function fitted to the energetics of bulk diamond, graphite, and a wide array of small hydrocarbons, such as C₂H₄, C₂H₆, C₆H₆, and the radicals \cdot CH₃ and \cdot C₂H₅.¹⁵ This potential has been successfully applied in MD simulations of a variety of chemical reactions,^{15,18,19} including hydrogen abstraction from hydrogen-terminated diamond surfaces.¹⁸ The Pt-C and Pt-H interactions are described by short-range Lennard-Jones 6/12 pair potentials,²⁰ which have been fitted to give bond distances and bond energies of 2.20 Å and 2.30 eV for Pt-C,^{21,22}

(10) (a) Harrison, D. E., Jr. *Crit. Rev. Solid State Mater. Sci.* **1988**, *14*, 51. (b) Garrison, B. J.; Winograd, N.; Harrison, D. E., Jr. *J. Chem. Phys.* **1978**, *69*, 1440.

(11) Garrison, B. J.; Winograd, N.; Deaven, D. M.; Reimann, C. T.; Lo, D. Y.; Tombrello, T. A.; Harrison, D. E., Jr.; Shapiro, M. H. *Phys. Rev. B* **1988**, *37*, 7197.

(12) Wucher, A.; Garrison, B. J. *J. Surf. Sci.* **1992**, *260*, 257.

(13) (a) Husinsky, W.; Nicolussi, G.; Betz, G. *Nucl. Instrum. Methods Phys. Res.* **1993**, *B82*, 323. (b) Wurz, P.; Husinsky, W.; Betz, G. *Appl. Phys.* **1991**, *A52*, 213.

(14) Urbassek, H. M. *Nucl. Instrum. Methods Phys. Res.* **1987**, *B18*, 587.

(15) (a) Brenner, D. W. *Phys. Rev. B* **1990**, *2*, 9458. (b) Brenner, D. W.; Harrison, J. A.; White, C. T.; Colton, R. J. *Thin Solid Films* **1991**, *206*, 220.

(16) Starke, U.; Barbieri, A.; Materer, N.; Van Hove, M. A.; Somorjai, G. A. *Surf. Sci.* **1993**, *286*, 1 and references therein.

(17) (a) Foiles, S. M.; Baskes, M. I.; Daw, M. S. *Phys. Rev.* **1986**, *33*, 7983. (b) Daw, M. S.; Baskes, M. I. *Phys. Rev. B* **1984**, *29*, 6443. (c) Daw, M. S.; Baskes, M. I. *Phys. Rev. Lett.* **1983**, *50*, 1285.

(18) (a) Brenner, D. W.; Robertson, D. H.; Carty, R. L.; Srivastava, D.; Garrison, B. J. *Mater. Res. Soc. Symp. Proc.* **1992**, *278*, 255. (b) Carty, R. L. Master's Thesis, Chemistry Department, The Pennsylvania State University, 1993.

(19) (a) Garrison, B. J.; Dawnkaski, E. J.; Srivastava, D.; Brenner, D. W. *Science* **1992**, *255*, 835. (b) Harrison, J. A.; White, C. T.; Colton, R. J.; Brenner, D. W. *Surf. Sci.* **1992**, *271*, 57. (c) Brenner, D. W.; Harrison, J. A. *Am. Ceram. Soc. Bull.* **1992**, *71*, 1821.

(20) Allen, M. P.; Tildesley, D. J. *Computer Simulations in Liquids*; Oxford University Press: Oxford, 1987; p 9.

(21) Carter, E. A.; Koel, B. E. *Surf. Sci.* **1990**, *226*, 339.

(22) Low, J. J.; Goddard, W. A. *J. Am. Chem. Soc.* **1986**, *108*, 6115.

(1) Recent reviews include the following: (a) Winograd, N. *Anal. Chem.* **1993**, *65*, 622a. (b) Benninghoven, A.; Hagenyhoff, B.; Niehuis, E. *Anal. Chem.* **1993**, *65*, 630a. (c) Winograd, N.; Garrison, B. J. In *Ion Spectroscopies for Surface Analysis*; Czanderna, A. W., Hercules, D. M., Eds.; Plenum Press: New York, 1991; pp 45-141. (d) Detter, L. D.; Hand, O. W.; Cooks, R. G.; Walton, R. A. *Mass Spectrom. Rev.* **1998**, *7*, 465. (e) Sunner, J.; Morales, A.; Kebarle, P. *Int. J. Mass Spectrom. Ion Processes* **1988**, *86*, 169. (f) Pachuta, S. J.; Cooks, R. G. *Chem. Rev.* **1987**, *87*, 647. (g) Benninghoven, A. *Springer Ser. Chem. Phys.* **1984**, *36*, 342. (h) Rinehart, K. L., Jr. *Science* **1982**, *218*, 254.

(2) Creighton, J. R.; White, J. M. *Surf. Sci.* **1983**, *129*, 327.

(3) Grade, H.; Winograd, N.; Cooks, R. G. *J. Am. Chem. Soc.* **1977**, *99*, 7725.

(4) Moon, D. W.; Bleiler, R. J.; Karwacki, E. J.; Winograd, N. *J. Am. Chem. Soc.* **1983**, *105*, 2916.

(5) (a) Garrison, B. J. *Int. J. Mass Spectrom. Ion Phys.* **1983**, *53*, 243. (b) Garrison, B. J. *J. Am. Chem. Soc.* **1982**, *104*, 6211. (c) Garrison, B. J. *J. Am. Chem. Soc.* **1980**, *102*, 6553.

(6) Lauderback, L. L.; Ang, M. L.; Murray, H. C. *J. Chem. Phys.* **1990**, *93*, 6041.

(7) Foley, K. E.; Winograd, N.; Garrison, B. J.; Harrison, D. E., Jr. *J. Chem. Phys.* **1984**, *80*, 5254.

(8) Williams, D. H.; Findeis, A. F.; Naylor, S.; Gibson, B. W. *J. Am. Chem. Soc.* **1987**, *109*, 1980.

(9) Brenner, D. W.; Garrison, B. J. *Phys. Rev. B* **1986**, *34*, 5782.




Figure 1. Two pathways found for H abstraction. The light gray, black, and white spheres represent the Pt, C, and H atoms, respectively. The H atoms that eventually form the H₂ molecule are marked with stars. In Scheme 1, the middle C₂H₃ is out of the plane of the reacting C₂H₃ molecules and does not enter into the H abstraction reaction. See the text for a description of the collision events.

and of 1.90 Å and 2.64 eV (in the 3-fold hollow) for Pt–H.^{23,24} The Ar–Pt, Ar–C, and Ar–H interactions are modeled by a screened Molière potential that allows only repulsive interactions.²⁵ A complete description of the calculational details including the merging of the Pt and C potentials and the hydrocarbon repulsive interactions will be given in full later.²⁶

Clusters varying in type from H₂ to Pt_xC_yH_z result from the collision events. In this communication, however, only the formation of H₂ will be highlighted. Unexpectedly, 66% of the H₂ predicted by these simulations is created via a hydrogen abstraction type mechanism. Although many variations exist, the two main abstraction mechanisms seen for the formation of H₂ are shown in Figure 1. In the predominant mechanism, Scheme 1, a H atom is emitted when the C₂H₃ molecule on the right is hit by either an energetic Pt atom, an Ar atom, or a CH_x (x = 0–3) fragment. Once emitted, this H atom is propelled in the direction of a H atom on an undisturbed C₂H₃ molecule. This results in the formation of a H–H bond and the breaking of a C–H bond. In the particular trajectory shown in Scheme 1, the H atom has 0.77 eV of kinetic energy when it enters (40 fs later) the neighbor shell of the undisturbed C₂H₃ molecule. Since the barrier to hydrogen abstraction for small hydrocarbons is between 0.2 and 0.45 eV,²⁷ sufficient energy is available to overcome this barrier. Scheme 2 shows an intriguing mechanism in which a C₂H₃ molecule is bumped such that one of its H atoms interacts with a H atom on the neighboring C₂H₃ molecule. H₂ is formed, and two C–H bonds are broken. The C–H bond on the

energetically excited C₂H₃ molecule is stretched to 1.37 Å, and the H atom has 0.98 eV of kinetic energy when it interacts with the neighboring C₂H₃ molecule. Each of the H abstraction reactions shown in Figure 1 is completed within 100 fs after the initial H atom is emitted; thus these are collisionally induced and not thermally induced reactions. Both of these pathways result in the same products: a free H₂ molecule and a C₂H₂ radical which remains adsorbed on the Pt surface. The H₂ molecules created are both vibrationally and rotationally hot, but out of the 90 formed, none have sufficient internal energy to dissociate. Although no experimental reports of H₂⁺ in the SIMS of C₂H₃ are available,² H₂⁺ has been observed in high-resolution time-of-flight SIMS images of polystyrene beads.²⁸

The identification of how molecular reactions proceed during keV particle bombardment should help simplify the complex relationship which exists between the structure of the molecular solid and the peaks observed in the mass spectra. In particular, several mechanisms have been ventured for the origin of the molecular ions, (M + H)⁺ and (M – H)⁻, as these peaks are often quite intense.^{1c-h,4,8} The “preformed ion” mechanism postulates that the (M ± H)[±] ions are created prior to the ion bombardment event in which they are ejected.^{1d,f} In the particular example of an organic film such as C₂H₃ adsorbed on a transition metal, it is not easy to imagine what chemical events would be necessary to give the adsorbed preformed (C₂H₃ + H)⁺ ion.²⁹ Perhaps instead, the formation of these ions proceeds by a mechanism analogous to that seen here for the formation of H₂. If the free H atom abstracts a H⁺ ion instead of a neutral H atom, then the (H + H)^o molecule can be seen as a neutral comparable to the (H + H)⁺ ion. Moreover, the abstraction process leaves an (M – H) entity, C₂H₂, on the surface which may later desorb. In addition to the “preformed ion” mechanism, reduction reactions of organic molecules have also been proposed to yield (M ± H)[±] ions.^{1d-f,8} These reduction reactions are proposed to proceed through either electron transfer alone or electron transfer with hydrogen attachment to the molecular ion.^{1d,8} Until now, it was believed that the H atoms emanated either from the substrate matrix, which in FAB is often glycerol, or from the decomposition of the molecular ion itself.^{1d,8} Although classical dynamics simulations do not take into account the presence of electrons, these simulations do suggest an alternate source of hydrogen.

Acknowledgment. The financial support of the National Science Foundation, the Office of Naval Research, and the IBM Selected University Research program is gratefully acknowledged. The Pennsylvania State University supplied a generous grant of computer time for these calculations. We also thank Nicholas Winograd, David E. Sanders, and Donald W. Brenner for insightful discussions.

(28) Brummel, C.; Winograd, N. Private communication.

(29) Although (C₂H₃ + H)⁺ is not seen experimentally, other molecular adsorbates on metal surfaces, such as C₅H₅N, do show a (M + H)⁺ peak in the SIMS spectra (see, for example, ref 4). C₅H₅N binds to the metal substrate through the N. Thus, for the preformed ion, (C₅H₅N + H)⁺, to exist, prior to any bombardment event, the metal–N bond must break, a H⁺–N bond must form, and this new C₅H₅NH⁺ species must remain adsorbed to the metal substrate until its ejection via an ensuing bombardment event.

(23) Sayers, C. M. *Surf. Sci.* **1984**, *143*, 411.

(24) Groppen, O.; Almlöf, J.; Wahlgren, J. *J. Chem. Phys.* **1992**, *96*, 8363.

(25) O'Connor, D. J.; Macdonald, R. J. *Radiat. Eff.* **1977**, *34*, 247.

(26) Taylor, R. S.; Garrison, B. J. Manuscript in preparation.

(27) Warnatz, J. In *Combustion Chemistry*; Gardiner, W. C., Ed.; Springer-Verlag: Berlin, 1984; p 271.